

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Thursday 15 October 2020

Afternoon (Time: 1 hour 30 minutes)

Paper Reference **9FM0/3A**

Further Mathematics

Advanced

Paper 3A: Further Pure Mathematics 1

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

--

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P62673A

©2020 Pearson Education Ltd.

1/1/1/1/1/1/



Pearson

1. Use l'Hospital's Rule to show that

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{(e^{\sin x} - \cos(3x) - e)}{\tan(2x)} = -\frac{3}{2}$$

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4.

$$f(x) = x^4 \sin(2x)$$

Use Leibnitz's theorem to show that the coefficient of $(x - \pi)^8$ in the Taylor series expansion of $f(x)$ about π is

$$\frac{a\pi + b\pi^3}{315}$$

where a and b are integers to be determined.

(8)

$$\left[\begin{array}{l} \text{The Taylor series expansion of } f(x) \text{ about } x = k \text{ is given by} \\ f(x) = f(k) + (x - k)f'(k) + \frac{(x - k)^2}{2!}f''(k) + \dots + \frac{(x - k)^r}{r!}f^{(r)}(k) + \dots \end{array} \right]$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



8.
$$f(x) = \frac{3}{13 + 6\sin x - 5\cos x}$$

Using the substitution $t = \tan\left(\frac{x}{2}\right)$

(a) show that $f(x)$ can be written in the form

$$\frac{3(1+t^2)}{2(3t+1)^2+6} \quad (3)$$

(b) Hence solve, for $0 < x < 2\pi$, the equation

$$f(x) = \frac{3}{7}$$

giving your answers to 2 decimal places where appropriate.

(5)

(c) Use the result of part (a) to show that

$$\int_{\frac{\pi}{3}}^{\frac{4\pi}{3}} f(x) dx = K \left(\arctan\left(\frac{\sqrt{3}-9}{3}\right) - \arctan\left(\frac{\sqrt{3}+3}{3}\right) + \pi \right)$$

where K is a constant to be determined.

(8)



