

## 9FM0/3A: Further Pure Mathematics 01 Mark scheme

Question	Scheme	Marks	AOs
1	$\frac{dx}{d\theta} = a \sec \theta \tan \theta, \frac{dy}{d\theta} = b \sec^2 \theta$	$k \frac{x}{a^2} - n \frac{y}{b^2} \frac{dy}{dx} = 0$ where $k > 0$ and $n > 0$	M1 2.1
	$\frac{dy}{dx} = \frac{b \sec^2 \theta}{a \sec \theta \tan \theta} \left( = \frac{b \sec \theta}{a \tan \theta} = \frac{b}{a \sin \theta} \right)$	$\frac{dy}{dx} = \frac{b^2 x}{a^2 y} = \frac{b \sec \theta}{a \tan \theta}$	A1 1.1b
	$y - b \tan \theta = \left( \frac{b \sec \theta}{a \tan \theta} \right) (x - a \sec \theta)$		M1 1.1b
	$y a \tan \theta = x b \sec \theta - a b^*$		A1* 2.1
		(4)	

(4 marks)

### Notes:

**M1:** Differentiates in an attempt to find  $\frac{dy}{dx}$   
 either differentiates  $x$  and  $y$  and divides  $\frac{dy}{d\theta}$  by  $\frac{dx}{d\theta}$   
 or achieves  $k \frac{x}{a^2} - n \frac{y}{b^2} \frac{dy}{dx} = 0$  where  $k > 0$  and  $n > 0$

**A1:** Correct expression for  $\frac{dy}{dx}$

**M1:** Uses  $y - b \tan \theta = \text{'their } \frac{dy}{dx} \text{'}$ ,  $(x - a \sec \theta)$

**A1\*:** Uses correct algebra and trig identities to achieve the correct equation of the tangent.

Question	Scheme	Marks	AOs
2(a)	$\mathbf{b} \times \mathbf{c} = 0\mathbf{i} + 5\mathbf{j} + 5\mathbf{k}$ or $\begin{pmatrix} 0 \\ 5 \\ 5 \end{pmatrix}$	M1	1.1b
	area of triangle $OBC = \frac{1}{2}  0\mathbf{i} + 5\mathbf{j} + 5\mathbf{k}  = \frac{1}{2} \sqrt{50} = \frac{5}{2} \sqrt{2}$ o.e.	A1	2.2a
		(2)	
(b)	$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 0 \\ 5 \\ 5 \end{pmatrix} = 0 + 5 + 0 = 5$	M1	1.1b
	volume of tetrahedron $OABC = \frac{1}{6} \times  5  = \frac{5}{6}$	A1	2.2a
		(2)	
<b>(4 marks)</b>			
<b>Notes:</b>			
<p>(a)</p> <p><b>M1:</b> Attempts the vector product <math>\mathbf{b} \times \mathbf{c}</math>, with at least two correct terms.</p> <p><b>A1:</b> Deduces area of triangle <math>OBC = \frac{1}{2}  \mathbf{b} \times \mathbf{c} </math></p>			
<p>(b)</p> <p><b>M1:</b> Attempt at the triple scalar product <math>\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})</math></p> <p><b>A1:</b> Deduces volume of tetrahedron <math>OABC = \frac{1}{6}  \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) </math></p>			

Question	Scheme	Marks	AOs
<b>3(a)</b>	$4\tan x + 3\cot\left(\frac{x}{2}\right) \sec^2\left(\frac{x}{2}\right)$ $= 4\tan x + \frac{3}{\tan\left(\frac{x}{2}\right)}\left(1 + \tan^2\left(\frac{x}{2}\right)\right)$ $= 4\left(\frac{2t}{1-t^2}\right) + \frac{3}{t}(1+t^2)$	M1	2.1
	$\left(\frac{8t}{1-t^2}\right) + \frac{3(1+t^2)}{t} = 0$ $8t^2 + 3(1+t^2)(1-t^2) = 0 \text{ or } \frac{8t^2+3(1+t^2)(1-t^2)}{t(1-t^2)} = 0$	M1	1.1b
	$3t^4 - 8t^2 - 3 = 0$ *	A1*	1.1b
		<b>(3)</b>	
<b>(b)</b>	Solves quadratic for $t^2$ by factorising, quadratic formula, calculator $(3t^2 + 1)(t^2 - 3) = 0, t^2 = \frac{8 \pm \sqrt{(-8)^2 - 4(3)(-3)}}{2(3)}$ $t^2 = 3, t^2 = -\frac{1}{3}$ leading to value for $t = \dots$	M1	3.1a
	$t = \pm\sqrt{3}$	A1	1.1b
	Finds two correct values for $x = -\frac{4\pi}{3}, -\frac{2\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$	M1	1.1b
	All correct values for $x = -\frac{4\pi}{3}, -\frac{2\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$	A1	2.2a
		<b>(4)</b>	
<b>(7 marks)</b>			
<b>Notes:</b>			
<b>(a)</b>			
<b>M1:</b> Expresses $3\cot\left(\frac{x}{2}\right) \sec^2\left(\frac{x}{2}\right)$ in terms of $\tan\left(\frac{x}{2}\right)$ and uses $t$ substitutions to obtain an expression in terms of $t$ only			
<b>M1:</b> Multiplies through by $t$ and $(1 - t^2)$ or forms a common denominator			
<b>A1*:</b> $3t^4 - 8t^2 - 3 = 0$ cso			
<b>(c)</b>			
<b>M1:</b> Solve the quadratic in $t^2$ leading to a value for $t$			
<b>A1:</b> Correct values for $t = \pm\sqrt{3}$			
<b>M1:</b> Finds two correct values for $x = -\frac{4\pi}{3}, -\frac{2\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$			
<b>A1:</b> All correct values for $x = -\frac{4\pi}{3}, -\frac{2\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$			

Question	Scheme	Marks	AOs
4	Solves $x^2 - 2x - 2 = 0$ or $x^2 + 2x - 2 = 0$	M1	1.1b
	Solves both $x^2 - 2x - 2 = 0$ and $x^2 + 2x - 2 = 0$	M1	3.1a
	$x = \frac{2 \pm \sqrt{12}}{2} = 1 \pm \sqrt{3}$ and $x = \frac{-2 \pm \sqrt{12}}{2} = -1 \pm \sqrt{3}$	A1	1.1b
	Deduces the required roots are $x = 1 + \sqrt{3}$ and $x = -1 + \sqrt{3}$	M1	2.2a
	e.g. $\{x \in \mathbb{R}: x < -1 + \sqrt{3}\} \cup \{x \in \mathbb{R}: x > 1 + \sqrt{3}\}$	A1	2.5
		(5)	
<b>(5 marks)</b>			
<b>Notes:</b>			
<p><b>M1:</b> Solves either <math>x^2 - 2x - 2 = 0</math> or <math>x^2 + 2x - 2 = 0</math></p> <p><b>M1:</b> Complete strategy to identify and solve all relevant equations and gets two critical values</p> <p><b>A1:</b> Correct exact values for <math>x</math>, may be unsimplified</p> <p><b>M1:</b> Deduces that the larger roots are required in each case</p> <p><b>A1:</b> Correct set of values given and correct set notation form</p>			

Question	Scheme	Marks	AOs
<b>5(i)</b>	<b><math>\mathbf{b} \times \mathbf{a}</math> is perpendicular to <math>\mathbf{a}</math> (and/or <math>\mathbf{b}</math>)</b>	M1	2.4
	Therefore <b><math>\mathbf{a} \cdot (\mathbf{b} \times \mathbf{a}) = 0</math></b>	A1	1.1b
		<b>(2)</b>	
<b>(ii)</b>	<b><math>\mathbf{a} \times \mathbf{b} = \mathbf{a} \times \mathbf{c} \Rightarrow \mathbf{a} \times (\mathbf{b} - \mathbf{c}) = \mathbf{0}</math></b>	M1	3.1a
	As <b><math>\mathbf{a} \neq \mathbf{0}</math></b> and <b><math>\mathbf{b} \neq \mathbf{c}</math></b> then <b><math>\mathbf{a}</math></b> is parallel to <b><math>(\mathbf{b} - \mathbf{c})</math></b> therefore <b><math>\mathbf{b} - \mathbf{c} = \lambda \mathbf{a}</math></b>	A1	2.4
		<b>(2)</b>	
<b>(4 marks)</b>			
<b>Notes:</b>			
<b>(i)</b> <b>M1:</b> Reasoning that <b><math>\mathbf{b} \times \mathbf{a}</math></b> is perpendicular to <b><math>\mathbf{a}</math></b> <b>A1:</b> <b><math>\mathbf{a} \cdot (\mathbf{b} \times \mathbf{a}) = 0</math></b>			
<b>(ii)</b> <b>M1:</b> Collecting on to one side and factorising <b>A1:</b> Reasoning: as <b><math>\mathbf{a} \neq \mathbf{0}</math></b> and <b><math>\mathbf{b} \neq \mathbf{c}</math></b> then <b><math>\mathbf{a}</math></b> is parallel to <b><math>(\mathbf{b} - \mathbf{c})</math></b> therefore <b><math>\mathbf{b} - \mathbf{c} = \lambda \mathbf{a}</math></b>			

Question	Scheme	Marks	AOs
<b>6(a)</b>	$x_0 = 0, \quad y_0 = 1, \quad \left(\frac{dy}{dx}\right)_0 = 0 - 1 = -1$	B1	1.1b
	$y_1 \approx y_0 + h \left(\frac{dy}{dx}\right)_0 = 1 + 0.05(-1) = \dots$	M1	1.1b
	$= 0.95$	A1	1.1b
	$\left(\frac{dy}{dx}\right)_1 = 0.05^2 - 0.95^2 = -0.9$	M1	2.1
	$y_2 \approx y_1 + h \left(\frac{dy}{dx}\right)_1 = 0.95 + 0.05(-0.9) = \dots$		
	$= 0.905$	A1	1.1b
	(5)		
<b>(b)</b>	$\frac{dy}{dx} = x^2 - y^2 \Rightarrow \frac{d^2y}{dx^2} = 2x - 2y \frac{dy}{dx}$	B1	2.1
	$\frac{d^3y}{dx^3} = 2 - \lambda y \frac{d^2y}{dx^2} \pm \mu \left(\frac{dy}{dx}\right)^2$ or substituting in $\frac{dy}{dx} = x^2 - y^2$ so that $\frac{d^2y}{dx^2} = 2x - 2yx^2 + 2y^3$ $\Rightarrow \frac{d^3y}{dx^3} = 2 \pm \alpha xy \pm \beta x^2 \frac{dy}{dx} + \delta y^2 \frac{dy}{dx}$	M1	1.1b
	$\frac{d^3y}{dx^3} = 2 - 2y \frac{d^2y}{dx^2} - 2 \left(\frac{dy}{dx}\right)^2$ or $\frac{d^3y}{dx^3} = 2 - 4xy - 2x^2 \frac{dy}{dx} + 6y^2 \frac{dy}{dx}$	A1	1.1b
		(3)	
<b>(c)</b>	$x = 0, y = 1 \Rightarrow \frac{dy}{dx} = -1$	M1	2.2a
	$\frac{d^2y}{dx^2} = 2(0) - 2(1)(-1) = 2$	M1	1.1b
	$\frac{d^3y}{dx^3} = 2 - 2(1)(2) - 2(-1)^2 = -4$	A1	1.1b
	$y = y(0) + y'(0)x + \frac{y''(0)x^2}{2} + \frac{y'''(0)x^3}{6} + \dots$	M1	2.5
	Series solution $y = 1 - x + x^2 - \frac{2}{3}x^3$	A1	1.1b
		(5)	
<b>(13 marks)</b>			

**Notes:****(a)**

**B1:**  $\left(\frac{dy}{dx}\right)_0 = -1$

**M1:** Applies the approximation formula with  $y_0$  and their value for  $\left(\frac{dy}{dx}\right)_0$ 

**A1:**  $y_1 = 0.95$

**M1:** Finds  $\left(\frac{dy}{dx}\right)_1$  and applies the approximation formula with their values for  $y_1$  and  $\left(\frac{dy}{dx}\right)_1$ 

**A1:**  $y_2 = 0.905$

**(b)**

**B1:** Differentiates to  $\frac{d^2y}{dx^2} = 2x - 2y \frac{dy}{dx}$

**M1:** Differentiates to the form  $\frac{d^3y}{dx^3} = 2 - \lambda y \frac{d^2y}{dx^2} \pm \mu \left(\frac{dy}{dx}\right)^2$  where  $\lambda > 0, \mu \neq 0$ or substituting in  $\frac{dy}{dx} = x^2 - y^2$  so that  $\frac{d^2y}{dx^2} = 2x - 2yx^2 + 2y^3$ 

$$\Rightarrow \frac{d^3y}{dx^3} = 2 \pm \alpha x \frac{dy}{dx} \pm \beta x^2 \frac{dy}{dx} + \delta y^2 \frac{dy}{dx} \text{ where } \delta > 0, \alpha, \beta \neq 0$$

**A1:**  $\frac{d^3y}{dx^3} = 2 - 2y \frac{d^2y}{dx^2} - 2 \left(\frac{dy}{dx}\right)^2$  or  $\frac{d^3y}{dx^3} = 2 - 4xy - 2x^2 \frac{dy}{dx} + 6y^2 \frac{dy}{dx}$

**(c)****B1:** Deduces the value for  $\frac{dy}{dx} = -1$ **M1:** Finds the values of  $\frac{d^2y}{dx^2}$  and  $\frac{d^3y}{dx^3}$ **A1:** Correct values for  $\frac{d^2y}{dx^2}$  and  $\frac{d^3y}{dx^3}$ **M1:** Substitutes into the correct formula and mathematical language, allow factorial notation**A1:** Correct series, must start with  $y = \dots$

Question	Scheme	Marks	AOs
7(a)	$A = 1000 \left(1 + \frac{5}{1200}\right)^{12} = 1051.16^*$	B1*	1.1b
		(1)	
(b)	Let $y = \left(1 + \frac{r}{100n}\right)^n$ so $\ln y = \ln \left(1 + \frac{r}{100n}\right)^n = n \ln \left(1 + \frac{r}{100n}\right)$	M1	3.1a
	$\lim_{n \rightarrow \infty} \ln y = \lim_{n \rightarrow \infty} n \ln \left(1 + \frac{r}{100n}\right) = \lim_{n \rightarrow \infty} \frac{\ln \left(1 + \frac{r}{100n}\right)}{1/n}$	M1	2.1
	$= \lim_{n \rightarrow \infty} \left[ \frac{-r/100n^2}{\left(1 + \frac{r}{100n}\right)} \div \frac{-1}{n^2} \right]$	dM1 A1	1.1b 1.1b
	$= \lim_{n \rightarrow \infty} \frac{r/100}{1 + \frac{r}{100n}} = \frac{r}{100}$	A1	1.1b
	$\lim_{n \rightarrow \infty} \left(1 + \frac{r}{100n}\right)^n = \lim_{n \rightarrow \infty} y = \lim_{n \rightarrow \infty} e^{\ln y} = e^{\lim_{n \rightarrow \infty} \ln y} = e^{\frac{r}{100}}$ *	A1*	2.1
		(6)	
(c)	$\lim_{n \rightarrow \infty} \left(1 + \frac{5}{100n}\right)^n = e^{0.05}$	B1	3.4
	Therefore $\lim_{n \rightarrow \infty} 1000 \left(1 + \frac{5}{100n}\right)^n = 1000e^{0.05}$	M1	2.2a
	Student has £1051.27 in their saving account after one year	A1	3.2a
		(3)	

**(10 marks)**

### Notes:

(a)

**B1\*:** Using  $P = 1000$ ,  $r = 5$  and  $n = 12$  to show  $A = 1051.16$

(b)

**M1:** Taking  $\ln$ 's to express  $\ln \left(1 + \frac{r}{100n}\right)^n = n \ln \left(1 + \frac{r}{100n}\right)$

**M1:** Expressing the limit as a quotient  $\lim_{n \rightarrow \infty} \ln y = \lim_{n \rightarrow \infty} \frac{\ln \left(1 + \frac{r}{100n}\right)}{1/n}$

**dM1:** Applies L'Hospital's rule and attempts to differentiate both the numerator and denominator.  
Depends on previous method mark

**A1:** Correct differentiation  $\lim_{n \rightarrow \infty} \left[ \frac{-r/100n^2}{\left(1 + \frac{r}{100n}\right)} \div \frac{-1}{n^2} \right]$  simplified or un-simplified

**A1:** Correct answer for the limit

**A1\*:** Fully correct proof with all mathematical notation cso

(c)

**B1:** Uses model and the result from part (b)  $\lim_{n \rightarrow \infty} \left(1 + \frac{0.05}{n}\right)^n = e^{0.05}$

**M1:** Deduces that the amount will be  $\lim_{n \rightarrow \infty} 1000 \left(1 + \frac{0.05}{n}\right)^n = 1000e^{0.05}$

**A1:** Give answer in pounds to 2 decimal places £1051.27

Question	Scheme	Marks	AOs
<b>8(a)</b>	$x = wt \Rightarrow \frac{dx}{dt} = \frac{dw}{dt} t + w$	B1	1.1b
	$t \left( \frac{dw}{dt} t + w \right) + 2t^2(wt) = wt(2t + 1)$	M1	2.1
	$t^2 \frac{dw}{dt} + tw + 2t^3w = 2wt^2 + wt$ $t^2 \frac{dw}{dt} + 2t^3w = 2wt^2$ Leading to $\frac{dw}{dt} + 2tw - 2w = 0$ *	A1*	1.1b
		(3)	
<b>(b)</b>	$\int \frac{1}{w} dw = 2 \int (1 - t) dt$	M1	3.1a
	$\ln w = 2t - t^2 (+c)$	M1	1.1b
	Uses correct exponential and ln work to reach $w = e^{2t - t^2 + c} = e^{2t - t^2} e^c = Ae^{2t - t^2}$	A1	2.1
	Displacement from $O$ is given by $x = wt = \dots$	M1	3.4
	$x = Ate^{2t - t^2}$ *	A1*	2.2a
		(5)	
<b>(c)</b>	Uses $x = 10$ when $t = 2$ to find the value of $A$ , $10 = 2A$ and achieves $x = 5te^{2t - t^2}$	B1	3.4
		(1)	
<b>(d)</b>	Sets $\frac{dx}{dt} = 0 \Rightarrow 2t^2x = x(2t + 1) \Rightarrow 2t^2 = 2t + 1$ or differentiates $x \Rightarrow \frac{dx}{dt} = 5e^{2t - t^2} + 5t(2 - 2t)e^{2t - t^2} = 0$ to form and solve a quadratic equation.	M1	3.1a
	$t = \frac{1 \pm \sqrt{3}}{2}$	A1	1.1b
	$x = 5 \left( \frac{1 + \sqrt{3}}{2} \right) e^{2 \left( \frac{1 + \sqrt{3}}{2} \right) - \left( \frac{1 + \sqrt{3}}{2} \right)^2}$	dM1	1.1b
	Maximum displacement = awrt 16.2 m or 162 cm	A1	3.2a
		(4)	
<b>(e)</b>	$x = 5te^{2t - t^2} = 5te^{2t} e^{-t^2}$ or $\frac{5te^{2t}}{e^{t^2}}$	M1	3.4
	As $t \rightarrow \infty$ , $e^{-t^2} \rightarrow 0$ or $e^{2t - t^2} \rightarrow 0 \therefore$ displacement from $O$ tends to 0 or $e^{t^2} \rightarrow \infty \therefore$ displacement from $O$ tends to 0	A1	2.4
		(2)	
<b>(15 marks)</b>			

**Notes:**

(a)

**B1:** Correct derivative  $\frac{dx}{dt}$

**M1:** Substitutes in their  $\frac{dx}{dt}$

**A1\*:** Completely correct proof

(b)

**M1:** Separates the variables correctly, with  $dw$  and  $dt$  the correct positions.

**M1:** Integrates both sides to the form  $\ln w = f(t)$  with or without  $+c$

**A1:** Uses correct exponential and  $\ln$  work to reach  $w = e^{2t-t^2+c} = e^{2t-t^2}e^c = Ae^{2t-t^2}$

Must have  $+c$  and a correct intermediate stage.

**M1:** Links their equation  $w = f(t)$  to the solution of the model equation correctly.

For  $x = t$  'their  $w$ '

**A1\*:** Deduces the correct general equation for the distance

(c)

**B1:** Uses  $x = 10$  when  $t = 2$  to find the correct value of  $A$

(d)

**M1:** Sets  $\frac{dx}{dt} = 0$  into the differential equation, or uses the product rule to differentiate  $x$  and sets

$\frac{dx}{dt} = 0$ , to form and solve a quadratic equation.

**A1:** Correct value(s) for  $t$

**dM1:** Dependent of previous method mark, substitutes their value of  $t$  to find a value for  $\underline{x}$ .

**A1:** Maximum displacement = awrt 16.2 m or 162 cm

(e)

**M1:** Using the model, separates the exponential terms

**A1:** Reason  $\therefore$  displacement from  $O$  tends to 0

Question	Scheme	Marks	AOs
<b>9(a)</b>	$\frac{a}{e} = \frac{8}{3}\sqrt{3}$	B1	3.4
	Uses $a = 2b$ and $b^2 = a^2(1 - e^2)$ Either $b^2 = 4b^2(1 - e^2)$ or $\frac{a^2}{4} = a^2(1 - e^2)$	M1	3.1a
	Either $\frac{1}{4} = 1 - e^2$ then $e = \frac{\sqrt{3}}{2}$ so $a = \left(\frac{\sqrt{3}}{2}\right)\left(\frac{8}{3}\sqrt{3}\right)$ or $e = \frac{3}{8\sqrt{3}} = \frac{a\sqrt{3}}{8}$ then $\frac{1}{4} = \left(1 - \frac{3a^2}{64}\right)$ leading to $a = \dots$	M1	2.1
	$a = 4$	A1	1.1b
	$b = \frac{a}{2} = 2$ and $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$	M1	1.1b
	$\frac{x^2}{16} + \frac{y^2}{4} = 1$	A1	1.1b
		(6)	
<b>(b)</b>	Foci $(\pm ae, 0)$ , $x = 4\left(\frac{\sqrt{3}}{2}\right) = 2\sqrt{3}$	M1	3.1b
	Water features at $(2\sqrt{3}, 0)$ and $(-2\sqrt{3}, 0)$	A1	3.4
		(2)	
<b>(c)</b>	Uses $PS = ePN$ , leading to $2 = \frac{\sqrt{3}}{2} PN$	M1	3.4
	$PN = \frac{4}{\sqrt{3}} = \frac{4}{3}\sqrt{3}$	A1	1.1b
	$x = \frac{8}{3}\sqrt{3} - \frac{4}{3}\sqrt{3} = \frac{4}{3}\sqrt{3}$	M1	1.1b
	$\frac{\left(\frac{4}{3}\sqrt{3}\right)^2}{16} + \frac{y^2}{4} = 1$ leading to a value for $y$	M1	1.1b
	Uses symmetry to find all 4 points $\left(\frac{4}{3}\sqrt{3}, \frac{2}{3}\sqrt{6}\right), \left(\frac{4}{3}\sqrt{3}, -\frac{2}{3}\sqrt{6}\right), \left(-\frac{4}{3}\sqrt{3}, \frac{2}{3}\sqrt{6}\right)$ and $\left(-\frac{4}{3}\sqrt{3}, -\frac{2}{3}\sqrt{6}\right)$	A1	3.4
		(5)	

	<b>Alternative 9(c)</b>		
	Solves $(x - '2\sqrt{3}')^2 + y^2 = 4$ or $(x + '2\sqrt{3}')^2 + y^2 = 4$ with the equation of their ellipse and follow through on their foci to find a value of $x$ or $y$	M1	3.4
	Any correct value of $x$ or $y$	A1	1.1b
	Uses symmetry to find another value of $x$ or $y$ or Solves $(x - '2\sqrt{3}')^2 + y^2 = 4$ and $(x + '2\sqrt{3}')^2 + y^2 = 4$ with the equation of their ellipse and follow through on their foci to find a value of $x$ or $y$	M1	1.1b
	Finds a complete point	M1	1.1b
	Finds all 4 points $(\frac{4}{3}\sqrt{3}, \frac{2}{3}\sqrt{6}), (\frac{4}{3}\sqrt{3}, -\frac{2}{3}\sqrt{6}), (-\frac{4}{3}\sqrt{3}, \frac{2}{3}\sqrt{6})$ and $(-\frac{4}{3}\sqrt{3}, -\frac{2}{3}\sqrt{6})$	A1	3.4
		(5)	

**(13 marks)**

**Notes:**

(a)

**B1:** Using half the length equals the  $x$  coordinate of the directrix

**M1:** Uses  $a = 2b$  and  $b^2 = a^2(1 - e^2)$

**M1:** For a complete method to find a value for  $a$

**A1:** For  $a = 4$

**M1:** Finding the value for  $b$  and substituting the values of  $a$  and  $b$  into the equation of an ellipse

**A1:** Correct equation of the ellipse, must square out  $a$  and  $b$ .

(b)

**M1:** For realising that the foci for the ellipse are required and finds  $x$ -coordinate of focus  $x = ae$

**A1:** Finds both coordinates for the water features

(c)

**M1:** Uses focus directrix property with  $PS = 2$  and their value for  $e$

**A1:** Correct distance for  $PN$

**M1:** Using  $x = \text{directrix} - PN$

**M1:** Substitutes value for  $x$  into their equation of the ellipse to find a value for  $y$

**A1:** All 4 correct points

(c) **Alternative**

**M1:** Solves simultaneously their equation of the ellipse and a circle with centre ('their foci', 0) and radius 2. Finds at least one value for  $x$  or  $y$

**A1:** A correct value of  $x = \pm \frac{4}{3}\sqrt{3}$  or  $y = \pm \frac{2}{3}\sqrt{6}$

**M1:** Or uses symmetry to find another values of  $x$  or  $y$ . Or solves simultaneously their equation of the ellipse and both circles with centre ('their foci', 0) and radius 2. Finds at least one value of  $x$  or  $y$ .

**M1:** Finds a complete coordinate

**A1:** All 4 correct points