

# Y2M8 XMQs and MS

(Total: 104 marks)

1. P3\_2018 Q6 . 6 marks - Y2M8 Further kinematics
2. P3\_2018 Q8 . 8 marks - Y2M8 Further kinematics
3. P3\_Sample Q8 . 10 marks - Y2M8 Further kinematics
4. P3\_Specimen Q6 . 4 marks - Y2M8 Further kinematics
5. P3\_Specimen Q7 . 7 marks - Y2M8 Further kinematics
6. P32\_2019 Q1 . 6 marks - Y2M8 Further kinematics
7. P32\_2019 Q2 . 8 marks - Y2M8 Further kinematics
8. P32\_2020 Q2 . 8 marks - Y2M8 Further kinematics
9. P32\_2020 Q3 . 12 marks - Y2M8 Further kinematics
10. P32\_2021 Q1 . 4 marks - Y2M8 Further kinematics
11. P32\_2021 Q5 . 14 marks - Y2M8 Further kinematics
12. P32\_2022 Q1 . 8 marks - Y2M8 Further kinematics
13. P32\_2022 Q3 . 9 marks - Y2M8 Further kinematics

## SECTION B: MECHANICS

Unless otherwise stated, whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ ms}^{-2}$  and give your answer to either 2 significant figures or 3 significant figures.

Answer ALL questions. Write your answers in the spaces provided.

6. At time  $t$  seconds, where  $t \geq 0$ , a particle  $P$  moves in the  $x$ - $y$  plane in such a way that its velocity  $\mathbf{v} \text{ ms}^{-1}$  is given by

$$\mathbf{v} = t^{-\frac{1}{2}}\mathbf{i} - 4t\mathbf{j}$$

When  $t = 1$ ,  $P$  is at the point  $A$  and when  $t = 4$ ,  $P$  is at the point  $B$ .

Find the exact distance  $AB$ .

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**Section B: MECHANICS**

| Question  | Scheme   | Marks | AOs       |
|---|--|-------|-----------|
| 6.  | Integrate $\mathbf{v}$ w.r.t. time   | M1    | 1.1a      |
|   | $\mathbf{r} = 2t^{\frac{1}{2}}\mathbf{i} - 2t^2\mathbf{j} (+ \mathbf{C})$  | A1    | 1.1b      |
|   | Substitute $t = 4$ and $t = 1$ into their $\mathbf{r}$   | M1    | 1.1b      |
|   | $t = 4, \mathbf{r} = 4\mathbf{i} - 32\mathbf{j} (+ \mathbf{C}); t = 1, \mathbf{r} = 2\mathbf{i} - 2\mathbf{j} (+ \mathbf{C})$ or $(4, -32); (2, -2)$ | A1    | 1.1b      |
|   | $\sqrt{2^2 + (-30)^2}$   | M1    | 1.1b      |
|   | $\sqrt{904} = 2\sqrt{226}$   | A1    | 1.1b      |
|   |  | (6)   |           |
|   |  |       | (6 marks) |
| <b>Notes: Allow column vectors throughout</b>   |  |       |           |
| <p><b>M1:</b> At least one power increasing by 1.</p> <p><b>A1:</b> Any correct (unsimplified) expression</p> <p><b>M1:</b> Must have attempted to integrate <math>\mathbf{v}</math>. Substitute <math>t = 4</math> and <math>t = 1</math> into their <math>\mathbf{r}</math> to produce 2 vectors (or 2 points if just working with coordinates).</p> <p><b>A1:</b> <math>4\mathbf{i} - 32\mathbf{j} (+ \mathbf{C})</math> and <math>2\mathbf{i} - 2\mathbf{j} (+ \mathbf{C})</math> or <math>(4, -32)</math> and <math>(2, -2)</math>. These can be seen or implied.</p> <p><b>M1:</b> Attempt at distance of form <math>\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}</math> for their points. Must have 2 non zero terms.</p> <p><b>A1:</b> <math>\sqrt{904} = 2\sqrt{226}</math> or any equivalent surd (exact answer needed)</p> |  |       |           |



| Question | Scheme  | Marks | AOs  |
|----------|---|-------|------|
| 8(a)     | Use of $\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$ : $(7\mathbf{i} - 10\mathbf{j}) = 2(2\mathbf{i} - 3\mathbf{j}) + \frac{1}{2}\mathbf{a}2^2$ | M1    | 3.1b |
|          | $\mathbf{a} = (1.5\mathbf{i} - 2\mathbf{j})$  | A1    | 1.1b |
|          | $ \mathbf{a}  = \sqrt{1.5^2 + (-2)^2}$  | M1    | 1.1b |
|          | $= 2.5 \text{ m s}^{-2}$ * GIVEN ANSWER   | A1*   | 2.1  |
|          |   | (4)   |      |
| (b)      | Use of $\mathbf{v} = \mathbf{u} + \mathbf{a}t = (2\mathbf{i} - 3\mathbf{j}) + 2(1.5\mathbf{i} - 2\mathbf{j})$   | M1    | 3.1b |
|          | $= (5\mathbf{i} - 7\mathbf{j})$   | A1    | 1.1b |
|          | $\mathbf{v} = (5\mathbf{i} - 7\mathbf{j}) + t(4\mathbf{i} + 8.8\mathbf{j}) = (5 + 4t)\mathbf{i} + (8.8t - 7)\mathbf{j}$ and<br>$(5 + 4t) = (8.8t - 7)$  | M1    | 3.1b |
|          | $t = 2.5 \text{ (s)}$   | A1    | 1.1b |
|          |   | (4)   |      |

(8 marks)

Notes: Allow column vectors throughout

(a)

**No credit for individual component calculations****M1:** Using a complete method to obtain the acceleration. **N.B.** Equation, in **a** only, could be obtained by two integrations**ALTERNATIVE****M1:** Use velocity at half-time ( $t = 1$ ) = Average velocity over time periodSo at  $t = 1$ ,  $\mathbf{v} = \frac{1}{2}(7\mathbf{i} - 10\mathbf{j})$  so  $\mathbf{a} = \frac{1}{2}(7\mathbf{i} - 10\mathbf{j}) - (2\mathbf{i} - 3\mathbf{j})$ **N.B.** could see  $(7\mathbf{i} - 10\mathbf{j}) = (4\mathbf{i} - 6\mathbf{j}) + 2\mathbf{a}$  as first line of working**A1:** Correct **a** vector**M1:** Attempt to find magnitude of their **a** using form  $\sqrt{a^2 + b^2}$ **A1\*:** Correct GIVEN ANSWER obtained correctly

(b)

**M1:** Using a complete method to obtain the velocity at A e.g. by use of  $\mathbf{v} = \mathbf{u} + \mathbf{a}t$  with  $t = 2$  and  $\mathbf{u} = 2\mathbf{i} - 3\mathbf{j}$  and their **a**OR: by use of  $\mathbf{s} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2$ OR: by integrating their **a**, with addition of  $\mathbf{C} = 2\mathbf{i} - 3\mathbf{j}$ , and putting  $t = 2$ **A1:** correct vector**M1:** Complete method to find equation in  $t$  only

e.g. by using  $\mathbf{v} = \mathbf{u} + \mathbf{a}t$ , with their  $\mathbf{u}$  and equating  $\mathbf{i}$  and  $\mathbf{j}$  components

---

**OR:** by integrating  $(4\mathbf{i} + 8.8\mathbf{j})$ , with addition of a constant, and equating  $\mathbf{i}$  and  $\mathbf{j}$  components.

**N.B.** Must be equating  $\mathbf{i}$  and  $\mathbf{j}$  components of a velocity vector and must be their velocity at A, to give an equation in  $t$  only for this M mark

**A1:** 2.5 (s)



| Question  | Scheme   | Marks      | AOs               |
|---|--|------------|-------------------|
| <b>8(a)</b>   | Use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$ : $(10.5\mathbf{i} - 0.9\mathbf{j}) = 0.6\mathbf{j} + 15\mathbf{a}$ | M1         | 3.1b              |
|   | $\mathbf{a} = (0.7\mathbf{i} - 0.1\mathbf{j}) \text{ m s}^{-2}$ Given answer                                       | A1         | 1.1b              |
|   |  | <b>(2)</b> |                   |
| <b>(b)</b>  | Use of $\mathbf{r} = \mathbf{ut} + \frac{1}{2} \mathbf{at}^2$  | M1         | 3.1b              |
|   | $\mathbf{r} = 0.6\mathbf{j}t + \frac{1}{2}(0.7\mathbf{i} - 0.1\mathbf{j})t^2$                                      | A1         | 1.1b              |
|   |  | <b>(2)</b> |                   |
| <b>(c)</b>  | Equating the <b>i</b> and <b>j</b> components of <b>r</b>  | M1         | 3.1b              |
|   | $\frac{1}{2} \leftarrow 0.7t^2 = 0.6t - \frac{1}{2} \leftarrow 0.1t^2$   | A1ft       | 1.1b              |
|   | $t = 1.5$  | A1         | 1.1b              |
|   |  | <b>(3)</b> |                   |
| <b>(d)</b>  | Use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$ : $\mathbf{v} = 0.6\mathbf{j} + (0.7\mathbf{i} - 0.1\mathbf{j})t$   | M1         | 3.1b              |
|   | Equating the <b>i</b> and <b>j</b> components of <b>v</b>  | M1         | 3.1b              |
|   | $t = 0.75$   | A1 ft      | 1.1b              |
|   |  | <b>(3)</b> |                   |
|   |  |            | <b>(10 marks)</b> |
| <b>Notes:</b>   |  |            |                   |
| <b>(a)</b>  |  |            |                   |
| <b>M1:</b> for use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$                               |  |            |                   |
| <b>A1:</b> for given answer correctly obtained  |  |            |                   |
| <b>(b)</b>  |  |            |                   |
| <b>M1:</b> for use of $\mathbf{r} = \mathbf{ut} + \frac{1}{2} \mathbf{at}^2$                |  |            |                   |
| <b>A1:</b> for a correct expression for <b>r</b> in terms of $t$                            |  |            |                   |
| <b>(c)</b>  |  |            |                   |
| <b>M1:</b> for equating the <b>i</b> and <b>j</b> components of their <b>r</b>              |  |            |                   |
| <b>A1ft:</b> for a correct equation following their <b>r</b>                                |  |            |                   |
| <b>A1:</b> for $t = 1.5$  |  |            |                   |
| <b>(d)</b>  |  |            |                   |
| <b>M1:</b> for use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$ for a general $t$             |  |            |                   |
| <b>M1:</b> for equating the <b>i</b> and <b>j</b> components of their <b>v</b>              |  |            |                   |
| <b>A1ft:</b> for $t = 0.75$ , or a correct follow through answer from an incorrect equation |  |            |                   |



**9MA0/03 Mock Paper: Part B Mechanics Mark scheme**

| Question   | Scheme   | Marks      | AOs  |
|--|--|------------|------|
| <b>1</b>   | $\mathbf{r} = (-4.5\mathbf{i} + 3\mathbf{j})$                                      | B1         | 1.1b |
|  | Use of $\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$                       | M1         | 3.1b |
|  | $(-4.5\mathbf{i} + 3\mathbf{j}) = 3\mathbf{u} + 0.5(\mathbf{i} - 2\mathbf{j}) 3^2$ | A1ft       | 1.1b |
|  | $\mathbf{u} = (-3\mathbf{i} + 4\mathbf{j})$  | A1         | 1.1b |
|  |  | <b>(4)</b> |      |
| <b>(4 marks)</b>   |  |            |      |
| <b>Notes:</b>  |  |            |      |
| <p><b>B1:</b> Correct displacement vector</p> <p><b>M1:</b> Use of correct strategy and/or formula to give equation in <math>\mathbf{u}</math> only (could be obtained by two integrations)</p> <p><b>A1ft:</b> Correct equation in <math>\mathbf{u}</math> only, following their displacement vector</p> <p><b>A1:</b> Correct answer</p> |  |            |      |

7. A particle,  $P$ , moves under the action of a single force in such a way that at time  $t$  seconds, where  $t \geq 0$ , its velocity  $\mathbf{v}$  m s<sup>-1</sup> is given by

$$\mathbf{v} = (t^2 - 3t) \mathbf{i} - 12t \mathbf{j}$$

The mass of  $P$  is 0.5 kg.

Find the time at which the magnitude of the force acting on  $P$  is 6.5 N.

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

| Question  | Scheme   | Marks      | AOs  |
|---|--|------------|------|
| 2   | Differentiate wrt $t$                              | M1         | 1.1a |
|   | $\mathbf{a} = (2t - 3) \mathbf{i} - 12 \mathbf{j}$ | A1         | 1.1b |
|   | $(2t - 3)^2 + (-12)^2$                             | M1         | 1.1b |
|   | $(2t - 3)^2 + (-12)^2 = (6.5 / 0.5)^2$ oe          | M1         | 2.1  |
|   | $4t^2 - 12t - 16 = 0$                              | A1         | 1.1b |
|   | $(t - 4)(t + 1) = 0$                               | M1         | 1.1b |
|   | $t = 4$  | A1         | 1.1b |
|   |  | <b>(7)</b> |      |
| <b>(7 marks)</b>  |  |            |      |
| <b>Notes:</b>   |  |            |      |
| <p><b>M1:</b> At least one power going down<br/> <b>A1:</b> A correct expression<br/> <b>M1:</b> Sum of squares of components (with or without square root) of <math>\mathbf{a}</math> or <math>\mathbf{F}</math><br/> <b>M1:</b> Equating magnitude to 6.5/0.5 or 6.5 as appropriate and squaring both sides<br/> <b>A1:</b> Correct quadratic = 0 in any form<br/> <b>M1:</b> Attempt to solve a 3 term quadratic<br/> <b>A1:</b> 4</p> |  |            |      |



| Question    | Scheme   | Marks  | AO   |
|-------------|--|--|------|
|             |  |  |      |
| <b>1(a)</b> | Differentiate $\mathbf{v}$   | M1   | 1.1a |
|             | $(\mathbf{a} =) 6\mathbf{i} - \frac{15}{2}t^{\frac{1}{2}}\mathbf{j}$   | A1   | 1.1b |
|             | $= 6\mathbf{i} - 15\mathbf{j} \text{ (m s}^{-2}\text{)}$   | A1   | 1.1b |
|             |  | (3)  |      |
| <b>1(b)</b> | Integrate $\mathbf{v}$   | M1   | 1.1a |
|             | $(\mathbf{r} =)(\mathbf{r}_0) + 3t^2\mathbf{i} - 2t^{\frac{5}{2}}\mathbf{j}$   | A1   | 1.1b |
|             | $= (-20\mathbf{i} + 20\mathbf{j}) + (48\mathbf{i} - 64\mathbf{j}) = 28\mathbf{i} - 44\mathbf{j} \text{ (m)}$                 | A1   | 2.2a |
|             |  | (3)  |      |
|             |  | (6)  |      |
| Marks       | Notes  |  |      |
|             | <b>N.B.</b> Accept column vectors throughout and condone missing brackets in working but they must be there in final answers |  |      |
| <b>1a</b>   | M1   | Use of $\mathbf{a} = \frac{d\mathbf{v}}{dt}$ with attempt to differentiate (both powers decreasing by 1)<br>M0 if $\mathbf{i}$ 's and $\mathbf{j}$ 's omitted and they don't recover |      |
|             | A1   | Correct differentiation in any form  |      |
|             | A1   | Correct and simplified.<br>Ignore subsequent working (ISW) if they go on and find the magnitude.   |      |
| <b>1b</b>   | M1   | Use of $\mathbf{r} = \int \mathbf{v} dt$ with attempt to integrate (both powers increasing by 1)<br>M0 if $\mathbf{i}$ 's and $\mathbf{j}$ 's omitted and they don't recover         |      |
|             | A1   | Correct integration in any form. Condone $\mathbf{r}_0$ not present  |      |
|             | A1   | Correct and simplified.  |      |



| Question | Scheme   | Marks  | AO   |
|----------|--|--|------|
| 2(a)     | $(\mathbf{v} = )\mathbf{C} + (2\mathbf{i} - 3\mathbf{j})t$   | M1   | 3.1a |
|          | $(\mathbf{v} = )(-\mathbf{i} + 4\mathbf{j}) + (2\mathbf{i} - 3\mathbf{j})t$  | A1   | 1.1b |
|          | $\frac{4 - 3T}{-1 + 2T} = \frac{-4}{3}$ oe   | M1   | 3.1a |
|          | $T = 8$  | A1   | 1.1b |
|          |  | (4)  |      |
| (b)      | $(\mathbf{s} = )\mathbf{C}t + (2\mathbf{i} - 3\mathbf{j})\frac{1}{2}t^2$ (+ D)   | M1   | 3.1a |
|          | $(\mathbf{s} = )(-\mathbf{i} + 4\mathbf{j})t + \frac{1}{2}(2\mathbf{i} - 3\mathbf{j})t^2$ (+ D)  | A1   | 1.1b |
|          | $AB = \sqrt{12^2 + 8^2}$<br><b>N.B. Beware you may see <math>4(2\mathbf{i} - 3\mathbf{j})</math> which leads to <math>\sqrt{(8^2 + 12^2)}</math> this is M0A0M0A0.</b> | M1   | 3.1a |
|          | $= 4\sqrt{13}$ (= 14.422051....) (m)   | A1cso  | 1.1b |
|          |  | (4)  |      |
|          |  | (8)  |      |
| Marks    | Notes  |  |      |
| 2a       | M1   | Use of $\mathbf{v} = \mathbf{u} + \mathbf{a}t$<br><b>OR</b> integration to give an expression of the form $\mathbf{C} + (2\mathbf{i} - 3\mathbf{j})t$ , where <b>C is a non-zero constant vector</b><br>M0 if <b>u</b> and <b>a</b> are reversed<br>Condone use of $\mathbf{a} = (2\mathbf{i} + 3\mathbf{j})$ for this M mark  |      |
|          | A1   | Any correct unsimplified expression seen or implied  |      |
|          | M1   | Correct use of ratios, <u>using a velocity vector</u> (must be using $\frac{-4}{3}$ ) to give equation <u>in T only</u><br>M0 if they equate $4 - 3T = -4$ and/or $-1 + 2T = 3$ and therefore M0 if they then divide to produce their equation   |      |
|          | A1   | Correct only   |      |
|          |  | <b>N.B.</b><br>(i) Can score the second M1A1 if they get $T = 8$ , using a calculator to solve two simultaneous equations, but if answer is wrong, and no equation in $T$ only, second M0<br>(ii) Can score M1A1 M1A1 if they get $T = 8$ , using trial and error, but if they don't get $T = 8$ , can only score max M1A1M0A0 |      |

|           |       |   |
|-----------|-------|---|
|           |       |   |
| <b>2b</b> | M1    | <p>Use of <math>\mathbf{s} = \mathbf{ut} + \frac{1}{2}\mathbf{at}^2</math> with <math>\mathbf{a} = (2\mathbf{i} - 3\mathbf{j})</math></p> <p><b>OR</b> integration to give an expression of the form <math>\mathbf{Ct} + (2\mathbf{i} - 3\mathbf{j})\frac{1}{2}t^2</math>, where <b>C</b> is <b>their non-zero constant <u>vector</u> from (a)</b></p> <p>Condone use of <math>\mathbf{a} = (2\mathbf{i} + 3\mathbf{j})</math> for this M mark</p> <p><b>OR</b> any other complete method using vector <b>suva</b>t equations</p> |
|           | A1    | Correct unsimplified expression seen or implied   |
|           | M1    | <p>Use of <math>t = 4</math> in their <b>s</b> (which must be a <b>displacement vector</b>) and then Pythagoras with the root sign</p> <p><b>N.B.</b> This M mark can be implied by a correct answer, otherwise we need to see Pythagoras used, with the root sign, for the M mark.</p>   |
|           | A1cso | Any surd form or 14 or better   |



| Question | Scheme  | Marks | AOs  |
|----------|---|-------|------|
| 2(a)     | Use of $\mathbf{v} = \mathbf{u} + \mathbf{a}t$ or integrate to give: $\mathbf{v} = (-2\mathbf{i} + 2\mathbf{j}) + 2(4\mathbf{i} - 5\mathbf{j})$   | M1    | 3.1a |
|          | $(6\mathbf{i} - 8\mathbf{j}) (\text{m s}^{-1})$   | A1    | 1.1b |
|          |   | (2)   |      |
| 2(b)     | Solve problem through use of $\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$ or integration<br>(M0 if $\mathbf{u} = \mathbf{0}$ )<br>Or any other complete method e.g use $\mathbf{v} = \mathbf{u} + \mathbf{a}T$ and $\mathbf{r} = \frac{(\mathbf{u} + \mathbf{v})T}{2}$ : | M1    | 3.1a |
|          | $-4.5\mathbf{j} = 2t\mathbf{j} - \frac{1}{2}t^2 5\mathbf{j}$ ( $\mathbf{j}$ terms only)   | A1    | 1.1b |
|          | The first two marks could be implied if they go straight to an algebraic equation.  |       |      |
|          | Attempt to equate $\mathbf{j}$ components to give equation in $T$ only<br>$(-4.5 = 2T - \frac{5}{2}T^2)$  | M1    | 2.1  |
|          | $T = 1.8$   | A1    | 1.1b |
|          |   | (4)   |      |
| 2(c)     | Solve problem by substituting <u>their</u> $T$ value (M0 if $T < 0$ ) into the $\mathbf{i}$ component equation to give an equation in $\lambda$ only:<br>$\lambda = -2T + \frac{1}{2}T^2 \times 4$  | M1    | 3.1a |
|          | $\lambda = 2.9$ or $2.88$ or $\frac{72}{25}$ oe   | A1    | 1.1b |
|          |   | (2)   |      |

**Notes: Accept column vectors throughout**

**(8 marks)**

|    |    |  |
|----|----|--|
| 2a | M1 | For any complete method to give a $\mathbf{v}$ expression with correct no. of terms with $t = 2$ used, so if integrating, must see the initial velocity as the constant.<br>Allow sign errors.   |
|    | A1 | Ca0 isw if they go on to find the speed.   |
| 2b | M1 | For any complete method to give a vector expression for $\mathbf{j}$ component of displacement in $t$ (or $T$ ) only, using $\mathbf{a} = (4\mathbf{i} - 5\mathbf{j})$ , so if integrating, RHS of equation must have the correct structure.<br>Allow sign errors. |
|    | A1 | Correct $\mathbf{j}$ vector equation in $t$ or $T$ . Ignore $\mathbf{i}$ terms.  |
|    | M1 | Must have earned 1 <sup>st</sup> M mark.   |

|    |    |   |
|----|----|---|
|    |    | Equate $\mathbf{j}$ components to give equation in $T$ (allow $t$ ) only (no $\mathbf{j}$ 's) which has come from a displacement. Equation must be a 3 term quadratic in $T$ .  |
|    | A1 | cao   |
| 2c | M1 | Must have earned 1 <sup>st</sup> M mark in (b)<br>Complete method - must have an equation in $\lambda$ only (no $\mathbf{i}$ 's) which has come from an appropriate displacement.. (e.g M0 if $\mathbf{a} = \mathbf{0}$ has been used)<br>Expression for $\lambda$ must be a quadratic in $T$ |
|    | A1 | cao   |



| Question                                       | Scheme   |   | Marks | AOs  |
|--|--|---|-------|------|
| <b>3(i)(a)</b>                                 | Integrate $\mathbf{a}$ wrt $t$ to obtain velocity  |   | M1    | 3.4  |
|  | $\mathbf{v} = (t - 2t^2)\mathbf{i} + \left(3t - \frac{1}{3}t^3\right)\mathbf{j} (+\mathbf{C})$ |   | A1    | 1.1b |
|  | $8\mathbf{i} - \frac{28}{3}\mathbf{j} \text{ (m s}^{-1}\text{)}$                               |   | A1    | 1.1b |
|  |  |   | (3)   |      |
| <b>3(i)(b)</b>                                 | Equate $\mathbf{i}$ component of $\mathbf{v}$ to zero  |   | M1    | 3.1a |
|  | $t - 2t^2 + 36 = 0$  |   | A1ft  | 1.1b |
|  | $t = 4.5$ (ignore an incorrect second solution)  |   | A1    | 1.1b |
|  |  |   | (3)   |      |
| <b>3(ii)</b>                                   | Differentiate $\mathbf{r}$ wrt to $t$ to obtain velocity                                       |   | M1    | 3.4  |
|  | $\mathbf{v} = (2t - 1)\mathbf{i} + 3\mathbf{j}$  |   | A1    | 1.1b |
|  | Use magnitude to give an equation in $t$ only  |   | M1    | 2.1  |
|  | $(2t - 1)^2 + 3^2 = 5^2$   |   | A1    | 1.1b |
|  | Solve problem by solving this equation for $t$   |   | M1    | 3.1a |
|  | $t = 2.5$  |   | A1    | 1.1b |
|  |  |   | (6)   |      |
| <b>(12 marks)</b>                              |  |   |       |      |
| <b>Notes: Accept column vectors throughout</b> |  |   |       |      |
| <b>3(i)(a)</b>                                 | M1   | At least 3 terms with powers increasing by 1 (but M0 if clearly just multiplying by $t$ )                                       |       |      |
|  | A1   | Correct expression  |       |      |
|  | A1   | Accept $8\mathbf{i} - 9.3\mathbf{j}$ or better. Isw if speed found.   |       |      |
| <b>3(i)(b)</b>                                 | M1   | Must have an equation in $t$ only (Must have integrated to find a velocity vector)  |       |      |
|  | A1ft   | Correct equation follow through on their $\mathbf{v}$ but must be a 3 term quadratic  |       |      |
|  | A1   | cao   |       |      |
| <b>3(ii)</b>                                   | M1   | At least 2 terms with powers decreasing by 1 (but M0 if clearly just dividing by $t$ )  |       |      |
|  | A1   | Correct expression  |       |      |
|  | M1   | Use magnitude to give an equation in $t$ only, must have differentiated to find a velocity (M0 if they use $\sqrt{x^2 - y^2}$ ) |       |      |

|  |    |   |
|--|----|---|
|  | A1 | Correct equation $\sqrt{(2t-1)^2 + 3^2} = 5$  |
|  | M1 | Solve a 3 term quadratic for $t$ which has come from differentiating and using a magnitude. This M mark can be implied by a correct answer with no working. |
|  | A1 | 2.5   |



| Question   | Scheme  |  | Marks | AOs  |
|--|---|--|-------|------|
| <b>1(a)</b>  | Use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$ with $t = 2$ : $\mathbf{v} = 4\mathbf{i} + 2(2\mathbf{i} - 3\mathbf{j})$       |  | M1    | 3.1a |
|  | <b>OR integration:</b> $\mathbf{v} = (2\mathbf{i} - 3\mathbf{j})t + 4\mathbf{i}$ , with $t = 2$                               |  |       |      |
|  | $\mathbf{v} = 8\mathbf{i} - 6\mathbf{j}$  |  | A1    | 1.1b |
|  |   |  | (2)   |      |
| <b>1(b)</b>  | Use of $\mathbf{r} = \mathbf{ut} + \frac{1}{2}\mathbf{at}^2$ at $t = 3$ :   |  | M1    | 3.1a |
|  | $(\mathbf{i} + \mathbf{j}) + \left[ 3 \times 4\mathbf{i} + \frac{1}{2} \times (2\mathbf{i} - 3\mathbf{j}) \times 3^2 \right]$ |  |       |      |
|  | <b>OR:</b> find $\mathbf{v}$ at $t = 3$ : $4\mathbf{i} + 3(2\mathbf{i} - 3\mathbf{j}) = (10\mathbf{i} - 9\mathbf{j})$         |  |       |      |
|  | then use $\mathbf{r} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$   |  |       |      |
| $(\mathbf{i} + \mathbf{j}) + \left[ \frac{1}{2} [4\mathbf{i} + (10\mathbf{i} - 9\mathbf{j})] \times 3 \right]$   |   |  |       |      |
| or $\mathbf{r} = \mathbf{vt} - \frac{1}{2}\mathbf{at}^2$   |   |  |       |      |
| $(\mathbf{i} + \mathbf{j}) + \left[ 3 \times (10\mathbf{i} - 9\mathbf{j}) - \frac{1}{2} \times (2\mathbf{i} - 3\mathbf{j}) \times 3^2 \right]$           |   |  |       |      |
| <b>OR integration:</b> $\mathbf{r} = (\mathbf{i} + \mathbf{j}) + \left[ (2\mathbf{i} - 3\mathbf{j}) \frac{1}{2}t^2 + 4\mathbf{i} \right]$ , with $t = 3$ |   |  |       |      |
| $\mathbf{r} = 22\mathbf{i} - 12.5\mathbf{j}$   |   | A1   | 2.2a  |      |
|  |   |  | (2)   |      |
| <b>(4 marks)</b>   |   |  |       |      |
| <b>Notes: Accept column vectors throughout</b>   |   |  |       |      |
| <b>1a</b>  | M1  | Complete method to find $\mathbf{v}$ , using $\mathbf{ruvat}$ or integration<br>(M0 if $\mathbf{i}$ and/or $\mathbf{j}$ is missing)  |       |      |
|  | A1  | Apply isw if they also find the speed  |       |      |
| <b>1b</b>  | M1  | Complete method to find the p.v. but this mark can be scored if they omit $(\mathbf{i} + \mathbf{j})$<br>i.e. the M1 is for the expression in the square bracket<br>If they integrate, the M1 is earned once the expression in the square bracket is seen<br>with $t = 3$<br>(M0 if $\mathbf{i}$ and/or $\mathbf{j}$ is missing) |       |      |
|  | A1  | cao  |       |      |



|    |    |                        |
|----|----|------------------------|
| 4c | B1 | B0 if incorrect extras |
|----|----|------------------------|

| Question          | Scheme   |   | Marks | AOs  |
|-------------------|--|---|-------|------|
|                   | <b>Allow column vectors throughout this question</b>   |   |       |      |
| <b>5(a)</b>       | Differentiate $\mathbf{v}$ wrt $t$   |   | M1    | 3.1a |
|                   | $\frac{3}{2}t^{-\frac{1}{2}}\mathbf{i} - 2\mathbf{j}$ isw  |   | A1    | 1.1b |
|                   |  |   | (2)   |      |
| <b>5(b)</b>       | $3t^{\frac{1}{2}} = 2t$  |   | M1    | 2.1  |
|                   | Solve for $t$  |   | DM1   | 1.1b |
|                   | $t = \frac{9}{4}$  |   | A1    | 1.1b |
|                   |  |   | (3)   |      |
| <b>5(c)</b>       | Integrate $\mathbf{v}$ wrt $t$   |   | M1    | 3.1a |
|                   | $\mathbf{r} = 2t^{\frac{3}{2}}\mathbf{i} - t^2\mathbf{j} (+\mathbf{C})$  |   | A1    | 1.1b |
|                   | $t = 1, \mathbf{r} = -\mathbf{j} \Rightarrow \mathbf{C} = -2\mathbf{i}$ so $\mathbf{r} = 2t^{\frac{3}{2}}\mathbf{i} - t^2\mathbf{j} - 2\mathbf{i}$ |   | A1    | 2.2a |
|                   |  |   | (3)   |      |
| <b>5(d)</b>       | $\sqrt{(3t^{\frac{1}{2}})^2 + (2t)^2} = 10$ or $(3t^{\frac{1}{2}})^2 + (2t)^2 = 10^2$  |   | M1    | 2.1  |
|                   | $9t + 4t^2 = 100$  |   | M(A)1 | 1.1b |
|                   | $t = 4$  |   | A1    | 1.1b |
|                   | $\mathbf{r} = 14\mathbf{i} - 16\mathbf{j}$   |   | M1    | 1.1b |
|                   | $\sqrt{14^2 + (-16)^2}$  |   | M1    | 3.1a |
|                   | $\sqrt{452} (2\sqrt{113})$ (m)   |   | A1    | 1.1b |
|                   |  |   | (6)   |      |
| <b>(14 marks)</b> |  |   |       |      |
| <b>Notes:</b>     |  |   |       |      |
| <b>5a</b>         | M1   | Both powers decreasing by 1 (M0 if vector(s) disappear but allow recovery)                  |       |      |
|                   | A1   | cao   |       |      |
| <b>5b</b>         | M1   | Complete method, using $\mathbf{v}$ , to obtain an equation in $t$ only, allow a sign error |       |      |
|                   | DM1  | Dependent on M1, solve for $t$  |       |      |

|           |       |  |
|-----------|-------|--|
|           | A1    | cao  |
| <b>5c</b> | M1    | Both powers increasing by 1 (M0 if vectors disappear but allow recovery) |
|           | A1    | Correct expression without <b>C</b>                                      |
|           | A1    | cao  |
| <b>5d</b> | M1    | Use of Pythagoras on <b>v</b> and 10 to set up equation in $t$           |
|           | M(A)1 | Correct 3 term quadratic in $t$  |
|           | A1    | cao  |
|           | M1    | Substitute their numerical $t$ value into their <b>r</b>                 |
|           | M1    | Use of Pythagoras to find the magnitude of their <b>r</b>                |
|           | A1    | cso  |



| Question   | Scheme   |   | Marks | AOs  |
|--|--|---|-------|------|
| 1(a)   | Put $t = 2$ in $\mathbf{v}$ and use Pythagoras: $\sqrt{12^2 + (-6\sqrt{2})^2}$                 |   | M1    | 3.1a |
|  | $\sqrt{216}, 6\sqrt{6}$ or 15 or better (m s <sup>-1</sup> )                                   |   | A1    | 1.1b |
|  |  |   | (2)   |      |
| 1(b)   | Differentiate $\mathbf{v}$ wrt $t$ to obtain $\mathbf{a}$                                      |   | M1    | 3.4  |
|  | $6t\mathbf{i} - 3t^{-\frac{1}{2}}\mathbf{j}$ oe (m s <sup>-2</sup> ) isw                       |   | A1    | 1.1b |
|  |  |   | (2)   |      |
| 1(c)   | Integrate $\mathbf{v}$ wrt $t$ to obtain $\mathbf{r}$  |   | M1    | 3.4  |
|  | $\mathbf{r} = t^3\mathbf{i} - 4t^{\frac{3}{2}}\mathbf{j} (+\mathbf{C})$                        |   | A1    | 1.1b |
|  | $(\mathbf{i} - 4\mathbf{j}) = 4^3\mathbf{i} - 4 \times 4^{\frac{3}{2}}\mathbf{j} + \mathbf{C}$ |   | M1    | 3.1a |
|  | $(-62\mathbf{i} + 24\mathbf{j})$ (m) isw e.g. if they go on to find the distance.              |   | A1    | 1.1b |
|  |  |   | (4)   |      |
| <b>(8 marks)</b>   |  |   |       |      |
| <b>Notes: Accept column vectors throughout apart from the answer to (b).</b> |  |   |       |      |
| 1a   | M1   | Need square root but -ve sign not required. Allow $\mathbf{i}$ 's and/or $\mathbf{j}$ 's to go missing from their $\mathbf{v}$ at $t = 2$ , provided they have applied Pythagoras correctly.  |       |      |
|  | A1   | cao<br><b>N.B.</b> Correct answer with no working can score 2 marks.  |       |      |
| 1b   | M1   | Both powers decreasing by 1. Allow a column vector.<br>M0 if $\mathbf{i}$ or $\mathbf{j}$ is missing but allow recovery in (b).   |       |      |
|  | A1   | cao. Do not accept a column vector.   |       |      |
| 1c   | M1   | Both powers increasing by 1<br>M0 if $\mathbf{i}$ or $\mathbf{j}$ is missing but allow recovery.  |       |      |
|  | A1   | $(\mathbf{r} = )$ not required  |       |      |
|  | M1   | Putting $\mathbf{r} = (\mathbf{i} - 4\mathbf{j})$ and $t = 4$ into their displacement <b>vector</b> expression which must have $\mathbf{C}$ (allow $C$ ) to give an equation in $\mathbf{C}$ only, seen or implied.<br>Must have attempted to integrate $\mathbf{v}$ for this mark to be available.<br><b>N.B.</b> $\mathbf{C}$ does not need to be found and <u>this is a method mark, so allow slips.</u> |       |      |
|  | A1   | cao   |       |      |



| Question                                       | Scheme  | Marks  | AOs  |
|--|---|--|------|
| <b>3(a)</b>                                    | $(4\mathbf{i} - \mathbf{j}) + (\lambda\mathbf{i} + \mu\mathbf{j}) = (4 + \lambda)\mathbf{i} + (-1 + \mu)\mathbf{j}$   | M1   | 3.4  |
|  | Use <b>ratios</b> to obtain an equation in $\lambda$ and $\mu$ <i>only</i>  | M1   | 2.1  |
|  | $\frac{(4 + \lambda)}{(-1 + \mu)} = \frac{3}{1}$ or $\frac{\frac{1}{4}(4 + \lambda)}{\frac{1}{4}(-1 + \mu)} = \frac{3}{1}$  | A1   | 1.1b |
|  | $\lambda - 3\mu + 7 = 0$ *      Allow $0 = \lambda - 3\mu + 7$ but nothing else.  | A1*  | 1.1b |
|  |   | <b>(4)</b>   |      |
| <b>(b)</b>                                     | $\lambda = 2 \Rightarrow \mu = 3$ ; Resultant force = $(6\mathbf{i} + 2\mathbf{j})$ (N)   | M1   | 3.1a |
|  | $(6\mathbf{i} + 2\mathbf{j}) = 4\mathbf{a}$ <b>OR</b> $ (6\mathbf{i} + 2\mathbf{j})  = 4a$  | M1   | 1.1b |
|  | Use of $\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2$ with $\mathbf{u} = \mathbf{0}$ , their $\mathbf{a}$ and $t = 4$ :<br>Or they may integrate their $\mathbf{a}$ twice with $\mathbf{u} = \mathbf{0}$ and put $t = 4$ : | DM1  | 2.1  |
|  | $\mathbf{r} = \frac{1}{2} \times \frac{(6\mathbf{i} + 2\mathbf{j})}{4} 4^2 = (12\mathbf{i} + 4\mathbf{j})$  |  |      |
|  | $\sqrt{12^2 + 4^2}$   | M1   | 1.1b |
|  | <b>ALTERNATIVE 1</b> for last two M marks:<br>Use of $s = ut + \frac{1}{2}at^2$ , with $u = 0$ , their $a$ and $t = 4$ :<br>$s = \frac{1}{2} \times \sqrt{1.5^2 + 0.5^2} \times 4^2$  | DM1  |      |
|  | Use of Pythagoras to find mag of $\mathbf{a}$ : $a = \sqrt{1.5^2 + 0.5^2}$  | M1   |      |
|  | <b>ALTERNATIVE 2</b> for last two M marks:<br>Use of $s = ut + \frac{1}{2}at^2$ , with $u = 0$ , their $a$ and $t = 4$ :<br>$s = \frac{1}{2} \times \left( \frac{\sqrt{6^2 + 2^2}}{4} \right) \times 4^2$                         | DM1  |      |
|  | Use of Pythagoras to find $ (6\mathbf{i} + 2\mathbf{j}) $ : $= \sqrt{6^2 + 2^2}$  | M1   |      |
|  | $\sqrt{160}$ , $2\sqrt{40}$ , $4\sqrt{10}$ oe or 13 or better (m)   | A1   | 1.1b |
|  | <b>(5)</b>  |  |      |
| <b>(9 marks)</b>                               |   |  |      |
| <b>Notes: Accept column vectors throughout</b> |   |  |      |
| <b>3a</b>                                      | M1  | Adding the two forces, $\mathbf{i}$ 's and $\mathbf{j}$ 's must be collected (or must be a <b>single</b> column vector) seen or implied                |      |
|  | M1  | Must be using ratios; Ignore an equation e.g. $(4 + \lambda)\mathbf{i} + (-1 + \mu)\mathbf{j} = 3\mathbf{i} + \mathbf{j}$ if they go on to use ratios. |      |

|           |         |   |
|-----------|---------|---|
|           |         | <p>However, if they write <math>4 + \lambda = 3</math> and <math>-1 + \mu = 1</math> then <math>3(-1 + \mu) = 3</math> so <math>4 + \lambda = 3(-1 + \mu)</math> with no use of a constant, it's M0</p> <p>They may use the acceleration, with a factor of <math>\frac{1}{4}</math> top and bottom, see alternative</p> <p><b>Allow one side of the equation to be inverted</b></p>   |
|           | A1      | Correct equation  |
|           | A1*     | Given answer correctly obtained. Must see at least one line of working, with the LH fraction 'removed'.   |
| <b>3b</b> | M1      | <p>Adding <math>\mathbf{F}_1</math> and <math>\mathbf{F}_2</math> to find the resultant force, <math>\lambda</math> and <math>\mu</math> must be substituted</p> <p><b>N.B.</b> M0 if they use <math>\mu = 2</math> coming from <math>-1 + \mu = 1</math> in part (a).</p>  |
|           | M1      | <p>Use of <math>\mathbf{F} = 4\mathbf{a}</math> Or <math> \mathbf{F}  = 4a</math>, where <math>\mathbf{F}</math> is <u>their</u> resultant. (including <math>3\mathbf{i} + \mathbf{j}</math>)</p> <p>This is an independent mark, so could be earned, for example, if they have subtracted the forces to find the 'resultant'</p> <p><b>N.B.</b> M0 if only using <math>\mathbf{F}_1</math> or <math>\mathbf{F}_2</math></p>  |
|           | DM<br>1 | <p>Dependent on previous M mark for</p> <p><b>Either:</b> use of <math>\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2</math> with <math>\mathbf{u} = \mathbf{0}</math>, their <math>\mathbf{a}</math> and <math>t = 4</math> to produce a displacement vector</p> <p><b>Or :</b> integrate twice, with <math>\mathbf{u} = \mathbf{0}</math>, their <math>\mathbf{a}</math> and <math>t = 4</math> to produce a displacement Vector</p> <p><b>Or:</b> use of <math>s = ut + \frac{1}{2}at^2</math> with <math>u = 0</math>, their <math>a</math> and <math>t = 4</math> to produce a length</p> |
|           | M1      | Use of Pythagoras, with square root, to find the magnitude of their displacement vector, $\mathbf{a}$ or $\mathbf{F}$ (M0 if only using $\mathbf{F}_1$ or $\mathbf{F}_2$ ) depending on which method they have used.  |
|           | A1      | cao   |